

**UNCLASSIFIED**  
**AD 410162**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY DDC 410162  
AS AD No.

AMRL-TDR-62-151

**SOME CARDIORESPIRATORY RESPONSES OF FLYING AND  
NON-FLYING PERSONNEL TO DIFFERENT VECTORS OF  
ACCELERATION WITH CORRELATION OF THESE RESPONSES  
TO OTHER VARIABLES**

ALVIN S. HYDE, M.D., PH.D.  
NEIL S. CHERNIACK, M.D.  
EVAN F. LINDBERG, CAPT., USAF, (MC)  
DOROTHY WHATELY

TECHNICAL DOCUMENTARY REPORT No. AMRL-TDR-62-151

DECEMBER 1962

BIOMEDICAL LABORATORY  
6570th AEROSPACE MEDICAL RESEARCH LABORATORIES  
AEROSPACE MEDICAL DIVISION  
AIR FORCE SYSTEMS COMMAND  
WRIGHT-PATTERSON AIR FORCE BASE, OHIO

## NOTICES

When US Government drawings, specifications, or other data are used for any purpose other than a definitely related government procurement operation, the government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Qualified requesters may obtain copies from ASTIA. Orders will be expedited if placed through the librarian or other person designated to request documents from ASTIA.

Do not return this copy. Retain or destroy.

Stock quantities available at Office of Technical Services, Department of Commerce, \$0.50.

### Change of Address

Organizations receiving reports via the 6570th Aerospace Medical Research Laboratories automatic mailing lists should submit the addressograph plate stamp on the report envelope or refer to the code number when corresponding about change of address.

## FOREWORD


This work was done by the Acceleration Section, Biophysics Branch, Biomedical Laboratory, 6570th Aerospace Medical Research Laboratories, Wright-Patterson Air Force Base, Ohio, under Project 7222, "Biophysics of Flight," and Task 722202, "Acceleration in Flight and Escape." The authors would like to thank Mr. John W. Frazier and Mr. Justin Taylor for their considerable technical assistance in this study.

## ABSTRACT

The cardiovascular and respiratory responses of test pilots are compared with the AMRL centrifuge panel members during headward (+  $G_z$ ) and forward (+  $G_x$ ) accelerations. Vital capacity decreased in all subjects with increasing forward acceleration. No significant difference existed between the cardiorespiratory performance of test pilots and that of the nonrated personnel that constitute the AMRL centrifuge panel. No correlation was noted between blackout and pulse rate, but correlation did exist between resting control and + 5 $G_z$  pulse rates. An extensive number of anthropometric measurements, indices of physical fitness, and measurements made during other stress did not correlate with tolerance to headward (+  $G_z$ ) acceleration or with respiratory performance during +  $G_z$  and +  $G_x$  acceleration.

## PUBLICATION REVIEW

This technical documentary report has been reviewed and is approved.

  
JOS. M. QUASHNOCK  
Colonel, USAF, MC  
Biomedical Laboratory

SOME CARDIORESPIRATORY RESPONSES OF FLYING  
AND NON-FLYING PERSONNEL TO DIFFERENT VECTORS OF ACCELERATION  
WITH CORRELATION OF THESE RESPONSES TO OTHER VARIABLES


INTRODUCTION

This report stems from a recent opportunity to study the responses to different profiles of acceleration of a select group of 31 test pilots. The motivation of this group was quite high, as shown by repeated psychiatric interviews and a battery of clinical psychologic tests (ref. 1). The subjects were, in fact, frankly competitive. Their responses to headward and forward accelerations (+  $G_z$  and +  $G_x$ ) are compared (except for +  $G_z$  tolerance) to those of the volunteer panel of the 6570th Aerospace Medical Research Laboratories (AMRL) human centrifuge in a series of 141 separate centrifugations. The comparative cardiorespiratory responses of both groups are discussed. In addition, attempted correlation of these findings with extensive anthropometric data, indices of "physical fitness" and other variables is presented.

METHODS

Accelerative force was created by a centrifuge with a 22-foot radius. Subjects were positioned in the free-swinging cab that maintains the vector resulting from the gravitational attraction of the earth and centripetal acceleration perpendicular to the floor of the cab. An unconventional restraint system (ref. 2) was used, formed of contoured tubular frame and raschel net (figure 1) with adjustments designed for anthropometric fit of the 5th to the 95th percentile of USAF personnel.

Three separate profiles of acceleration were studied (figure 2). The sequence of presentation of these profiles was varied randomly. Although anti-G suits were not used, the subjects were permitted whatever straining maneuvers they were accustomed to use in order to prevent or delay blackout.



1. Net and forced tubular restraint
2. Balanced benedict-roth spirometer
3. Linear potentiometer
4. Kirchhoff valve
5. Smooth-bore, semi-rigid 1 inch I.D. tubing
6. Center light task

Figure 1. Subject in Position on Free-Swinging Gondola of AMRL Human Centrifuge




Figure 2. Acceleration Profiles



During headward acceleration ( $+G_z$ ) the subjects were exactly parallel to the inertial force ( $0^\circ$ ) with thighs flexed  $90^\circ$  on the trunk and knees flexed  $90^\circ$ . Subjects in the lowest percentile group required the use of urethane cushions to prevent excessive footward sink during acceleration. An arbitrary limit of headward acceleration was set at  $+9 G_z$  since the margin of difference between the levels of blackout and unconsciousness becomes small above  $+8 G_z$ . If blackout did not occur by  $+9 G_z$  the centrifuge stopped automatically. Blackout was recognized by failure to respond within 2 seconds to a central light signal offered at least every 4 seconds by an experienced central observer. The largest error possible therefore (assuming that blackout occurred immediately after answering the center-light) would be 6 seconds and would represent 0.40 G.

Two test series were run in forward accelerations ( $+G_x$ ). In one the subjects were perpendicular to the inertial force and in the other they were inclined  $12^\circ$  toward the direction of acceleration (figure 2).

The profiles of accelerations seen in figure 2 define the plateaus during which vital capacity and 0.5 second timed expiratory capacity were obtained. The control values for these maneuvers were established with the subject in position on the centrifuge some 3 minutes prior to centrifugation.

The expired air for these pulmonary function tests was collected in a specially balanced Benedict-Roth respirometer from which the carbon dioxide absorbent chamber had been removed. Displacement of the respirometer bell was transmitted to a multichannel recorder by a linear potentiometer coupled to the pulley of the respirometer bell chain. Respiratory rate was monitored by a mouthpiece thermistor just proximal to a Kirchoff valve (25-cc dead space). This tubing circuitry is shown in figure 1. Semi-rigid smooth-rubber tubing of 1 inch inside diameter was used throughout the expiratory circuit. The volume of this tubing was approximately 750 cc.

Electrocardiographic tracings were obtained from a modification of placement of the standard lead II and were both monitored and recorded continuously during each run. At the higher accelerations, above 8-10 G, myographic noise from the tense skeletal muscle mass caused interference that obscured the baseline of the electrocardiograms. Unfortunately the mechanisms of the arrhythmias which occurred at the higher G levels were masked by this interference, since at 10-12 G the P wave and even T waves were difficult to read with reasonable certainty.

## RESULTS AND DISCUSSION

### Headward Acceleration

Table I shows the response to headward acceleration of gradual rate of onset (1 G/15 seconds) in terms of blackout and pulse rates. Because gradual onset of acceleration, as used in this study, allows ample time for cardiovascular reflexes to compensate for the inertial insult to circulation (ref. 3), the mean blackout,  $+7.0 G_z \pm 1.2$ , was considerably higher than that usually found in  $+G_z$  acceleration. The range for blackout in the test pilot group extended from  $+4.7$  to  $+9.0 G_z$ . Two of the test pilots reached this arbitrary cut-off point of  $+9.0 G_z$  without blacking out; a feat made more remarkable when one considers that anti-G suits were not used and the gradual onset of acceleration plus the vital capacity measurement kept these subjects above  $+5 G_z$  for 75 seconds of the 135 seconds required to reach  $+9 G_z$ . Tolerance to headward acceleration ( $+G_z$ ) was determined under different conditions in the centrifuge panel, so that comparison should not be made between the two groups. (The panel members were instructed to relax during acceleration; the test pilots were allowed to strain.)

TABLE I  
RESPONSE TO HEADWARD ACCELERATION (+ G<sub>Z</sub>) OF GRADUAL  
RATE OF ONSET (1 G/15 SECONDS) IN TERMS OF  
BLACKOUT AND PULSE RATES

Panel (N = 15) Pilots (N = 31)

	+ G <sub>Z</sub> (Headward Acceleration)				+ G <sub>X</sub> (Forward Acceleration)							
	Blackout (G)	Pre-run Control Pulse	5G Pulse (Beats/min)	Pulse at Blackout	0° Back Angle			12° Back Angle				
					Pre-run Control Pulse	5G Pulse	8G Pulse	Pre-run Control Pulse	5G Pulse	8G Pulse	12G Pulse	
Mean	7.0	104	160	175	87	108	107	87	111	111	118	
S. D.	1.2	18	24	21	16	14	15	14	17	15	21	
S. E.	0.2	3	6	4	3	3	3	2	3	3	4	
Mean	5.5	109	166	174	99	116	131	93	121	125	121	
S. D.	0.9	14	16	15	17	19	21	17	18	16	22	
S. E.	0.2	4	4	4	5	6	7	5	5	5	8	

S. D. = Standard Deviation

S. E. = Standard Error

N = Number of Subjects

No significant correlation was found to exist between the control pulse rate and the blackout level, the control pulse rate and the blackout pulse rate, or pulse rate at blackout and the blackout level. There was a real ( $0.01 > P > 0.001$ ) but moderate correlation ( $r = + 0.59$ ) existing between the control pulse rate and the pulse rate at +5 G<sub>Z</sub>. The mean increase from control to +5 G<sub>Z</sub> pulse rate was  $35 \pm 3$  percent.

Brief, relative bradycardia associated with one or more extrasystoles occurred near peak acceleration (viz: blackout level) in 6 of 31 members of the test pilot group. One of the 15 centrifuge panel members had a single ectopic beat (not associated with a bradycardia) and a second member had intermittent sinus arrest. The incidence of arrhythmias resulting from + G<sub>Z</sub> accelerations found in this study may be low since the technical quality of the electrocardiograms at high accelerations occasionally did not permit detection of the P wave or the P-R interval. In addition, the arrhythmias reported here may be related to the unusually prolonged accelerations and to the slow rate of onset used.

Table II shows the effect of headward acceleration (+ G<sub>Z</sub>) upon the vital capacity and timed expiratory capacity. At +5 G<sub>Z</sub> the vital capacity is 55 percent of the control value for the test pilot group and 48 percent in the centrifuge panel group. The control 0.5-second forced expiratory capacity was  $59 \pm 13$  percent for the pilots and  $49 \pm 14$  percent of the control vital capacity for the panel members, but at +5 G<sub>Z</sub> there was no significant changes between either group ( $63 \pm 19$  percent and  $64 \pm 17$  percent).

TABLE II  
COMPARISON OF CARDIORESPIRATORY RESPONSES  
OF PILOT AND PANEL GROUPS

		+ G <sub>Z</sub>		+ G <sub>X</sub>				
				0° Back Angle		12° Back Angle		
			5 G	5 G	8 G	5 G	8 G	12 G
PILOTS (N = 31)	Control Vital Capacity (Liters)	$\bar{x}$	4.00	3.75	3.75	3.70	3.70	3.70
		± SD	.48	0.45	0.45	0.48	0.48	0.48
	Experimental Vital Capacity (Liters)	$\bar{x}$	2.20	1.68	0.48	1.60	0.55	0.20
		± SD	0.41	0.43	0.37	0.53	0.43	0.22
	Control Forced Expiratory Capacity (0.5 sec., percent)	$\bar{x}$	59	59	59	55	55	55
		± SD	13	11	11	12	12	12
	Experimental Forced Expiratory Capacity (0.5 sec., percent)	$\bar{x}$	63	80	86	77	91	94
		± SD	19	18	26	20	13	13
PANEL (N = 15)	Control Vital Capacity (Liters)	$\bar{x}$	3.88	3.85	3.85	3.75	3.75	3.75
		± SD	0.73	0.58	0.58	0.57	0.57	0.57
	Experimental Vital Capacity (Liters)	$\bar{x}$	1.85	1.45	0.48	1.45	0.45	0.05
		± SD	0.68	0.33	0.45	0.43	0.40	0.08
	Control Forced Expiratory Capacity (0.5 sec., percent)	$\bar{x}$	49	50	50	53	53	53
		± SD	14	8	8	17	17	17
	Experimental Forced Expiratory Capacity (0.5 sec., percent)	$\bar{x}$	64	82	91	77	93	100
		± SD	17	16	24	16	3	0

 $\bar{x}$  = Mean

SD = Standard Deviation

### Forward Accelerations (+ $G_x$ )

Forward accelerations up to +12  $G_x$  did not seem to present a serious cardiovascular challenge. Table I shows that only moderate increase over the resting (control) pulse rates occurred during forward acceleration. No clinically significant difference was found between the pilot group and the panel group, although the panel members tended toward consistently higher pulse rates in all situations, including the prerun "resting" control.

One of the test pilots had four extrasystoles and another had sudden bradycardia; all of this occurred at peak (+12  $G_x$ ) accelerations.

Dermksian and Lamb (ref. 4) have repeatedly emphasized the significance of the well established causal relationship existent between respiratory maneuvers and cardiac arrhythmias. Since all subjects vigorously performed forced expiratory capacities two or more times during each profile of forward (+  $G_x$ ) and headward (+  $G_z$ ) acceleration, the relative lack of arrhythmias during these tests appeared a more provocative finding than their rare presence during this series. Fear of the vago-inhibitory reflexes previously held by the medical monitors at the AMRL centrifuge facility has been considerably attenuated as a result of this series as well as subsequent studies of acceleration characterized by repeated respiratory maneuvers (refs. 5, 6).

During forward acceleration, we are most interested in pulmonary responses since the primary limitation to +  $G_x$  (forward acceleration) is generally conceded to be respiratory distress of one sort or another. This distress is variable, and ranges from dyspnea through severe and limiting pain at various thoracic sites, characteristically being most intense during the inspiratory phase. No real difference in ventilatory capability during acceleration existed between the test panel used for such studies and the group of test pilots. Both groups showed comparable, although severe, reductions in vital capacity which was proportionately related to the magnitude of the accelerations (table II).

Also of interest was the lack of relationship found between the back angles of 0° and 12° and the volumes of air that could be moved during acceleration. Previously, a 25° back angle has been shown to strongly influence the rate of decrement of vital capacity during acceleration when compared with a 0° back angle (ref. 3). The present study showed that 12° does not influence vital capacity during forward accelerations up to +8  $G_x$  when compared to 0° since no significant change was relatable to this variable (table II).


### Correlation of Acceleration with other Variables

Because an unusually extensive anthropometric survey was made to characterize the test pilot group, and because their responses to other stress tests were concurrently studied, we expected that some anthropometric or other variable obtained without centrifugation might correlate with pulmonary performance and tolerance to both headward (+  $G_z$ ) and forward (+  $G_x$ ) acceleration, thus providing selective criteria. For example, one might reasonably expect tolerance to +  $G_z$  to relate directly to the difference in heart-to-eye height in a group of pilots of similar experience and proficiency. Regrettably, for prediction of performance would be a true convenience to all military surgeons, this was not found to occur.

Although 104 variables have been cross correlated for this group of 31 test pilots (ref. 1), only those correlations which are significant statistically ( $r = 0.45$  when  $N = 31$ ), or which bear interest by their lack of statistical significance will be presented. The following relationships are considered especially noteworthy and are charted in table III.

TABLE III  
CORRELATION BETWEEN + G<sub>x</sub> AND + G<sub>z</sub>  
WITH VARIOUS INDICES

NOTE: ● connects variables with correlation coefficient ( $r$ ) greater than +0.449, (or with  $N = 31$ , there is 99 percent confidence that this is not chance alone)



Of all the tests performed at other institutions or during exposure to other stresses, correlation existed primarily between vital capacity obtained during +  $G_X$  acceleration and vital capacity as established "on the ground" at the Lovelace Foundation. Vital capacity at +5  $G_Z$  did not correlate with vital capacity at ground, but did correlate with lean body mass as determined at the Lovelace Foundation.

Hunter (ref. 8) also has unsuccessfully attempted to correlate what he termed "heart-brain distance" and sitting height against threshold to headward (+  $G_Z$ ) acceleration. He used a rate of onset of 1 G/10 seconds; this report used 1 G/15 seconds.

Kennedy et al (ref. 9), in a similar study that used 100 aircrew members as subjects, found no relation existent between blackout threshold and age, weight, body measurement, and response to tilt tests.

Because pulmonary responses to +  $G_X$  and +  $G_Z$  accelerations and pulse and blackout response to +  $G_Z$  accelerations bore no relationship at all to a variety of anthropometric indices, hemodynamic responses to heat stress, as well as most "physical fitness" tests that have ever been construed, we concluded that cardiopulmonary tolerance to acceleration can be determined only by actual measurement during such accelerations. Further, we all believe that the only selective value to such test accelerations as performed with the human centrifuge is that of negative selection. That is, if a subject is hemodynamically susceptible to acceleration (as would be manifest by a very low threshold to blackout or by acceleration induced serious cardiac arrhythmias) such susceptibility could be noted. However, the presence of reduced tolerance to acceleration is improbable in experienced test pilots whose very survival represents proof of relative hemodynamic integrity.

#### SUMMARY

A group of experienced test pilots and a group of nonrated centrifuge panel members underwent headward (+  $G_Z$ ) and forward (+  $G_X$ ) accelerations on a contoured net seat. No significant difference was found to exist in the responses of these two groups.

Two of the members of the test pilot group were able to reach +9.0  $G_Z$  acceleration without the occurrence of blackout.

No correlation was found between pulse rate and blackout level. Correlation did exist between resting pulse rate and pulse rate at +5  $G_Z$ .

Vital capacity decreased about 50 percent in both groups at +5  $G_Z$ .

Forward accelerations up to +12  $G_X$  caused little change in pulse rate and relatively few, minor, EKG abnormalities despite the concurrent performance of vigorous respiratory maneuvers.

Vital capacity decreased with increasing forward acceleration. A change in back angle from 0° to 12° did not significantly influence the rate of decrement in vital capacity. Half second forced expiratory capacity represented an increasing portion of total vital capacity as the forward acceleration increased.

Blackout tolerance during headward acceleration (+  $G_Z$ ) and respiratory performance during headward (+  $G_Z$ ) and forward acceleration (+  $G_X$ ) did not correlate with an extensive number (104) of anthropometric measurements and physical fitness tests nor was correlation found between +  $G_X$  and +  $G_Z$  acceleration responses and measurements made during other stressful situations.

## REFERENCES

1. Wilson, C.L. (Editor), Project Mercury Candidate Evaluation Program, WADC Technical Report 59-505, Wright Air Development Center, Wright-Patterson Air Force Base, Ohio, December 1959.
2. Bowring, J.I.R., Proceedings of Institute of Environmental Sciences, National Meeting, pp 262-274, 1960.
3. Edelberg, R., J.P. Henry, J.A. Maciolek, E.W. Salzman, and G.D. Zuidema, "Comparison of Human Tolerance to Accelerations of Slow and Rapid Onset," J. Aviat. Med., Vol 27, pp 482-489, December 1956.
4. Dermksian, G., and L. Lamb, Cardiac Arrhythmias in Experimental Syncope, Report 59-16, Air University, School of Aviation Medicine, Randolph Air Force Base, Texas, November 1958.
5. Cherniack, N.S., A.S. Hyde, and F.W. Zechman, "The Effect of Transverse Acceleration on Pulmonary Function," J. Appl. Physiol., Vol 14, No. 6, pp 914-916, 1959.
6. Zechman, F.W., N.S. Cherniack, and A.S. Hyde, Ventilatory Response to Forward Acceleration, WADC Technical Report 59-584, Wright Air Development Center, Wright-Patterson Air Force Base, Ohio, September 1959.
7. Zechman, F.W., The Effect of Forward Acceleration on Vital Capacity, WADC-Technical Note 58-376, Wright Air Development Center, Wright-Patterson Air Force Base, Ohio, December 1958.
8. Hunter, S., Correlation of Heart-Brain Distance and of Sitting Height Against Acceleration Thresholds, FPRC No. 1048, Flying Personnel Research Committee, Great Britain Air Air Ministry, Farnborough, Hants, England, April 1958.
9. Kennedy, W.A., W.K. Kerr, W.R. Martin, B. Rose, and W.R. Frank, "Relation of Blackout Threshold to Age, Weight, Body Measurement and Cardiovascular Tests," Proceedings of 13th Meeting, Associate Committee on Aviation Medicine Research, National Research Council of Canada, Ottawa, February 1944.

<p>Aerospace Medical Division, 6570th Aerospace Medical Research Laboratories, Wright-Patterson AFB, Ohio Rpt No. AMRL-TDR-62-151. SOME CARDIO- RESPIRATORY RESPONSES OF FLYING AND NON-FLYING PERSONNEL TO DIFFERENT VECTORS OF ACCELERATION WITH CORRE- LATION OF THESE RESPONSES TO OTHER VARIABLES. Final report, Dec 62, iii + 9 pp., incl. illus., tables, 9 refs. Unclassified report</p> <p>The cardiovascular and respiratory responses of test pilots are compared with the AMRL cen- trifuge panel members during headward (+ G<sub>Z</sub>) and forward (+ G<sub>X</sub>) accelerations. Vital capacity decreased in all subjects with increasing for- ward acceleration. No significant</p> <p>( over )</p>	<p>UNCLASSIFIED</p> <ol style="list-style-type: none"> <li>1. Biophysics</li> <li>2. Circulatory responses</li> <li>3. Acceleration</li> <li>4. Mechanics</li> <li>5. Stress Tests</li> <li>6. Anthropology</li> <li>7. Selection</li> </ol> <p>I. AFSC Project 7222, Task 722202</p> <p>II. Biomedical Labora- tory</p> <p>UNCLASSIFIED</p>	<p>UNCLASSIFIED</p> <ol style="list-style-type: none"> <li>1. Biophysics</li> <li>2. Circulatory responses</li> <li>3. Acceleration</li> <li>4. Mechanics</li> <li>5. Stress Tests</li> <li>6. Anthropology</li> <li>7. Selection</li> </ol> <p>I. AFSC Project 7222, Task 722202</p> <p>II. Biomedical Labora- tory</p> <p>UNCLASSIFIED</p>	<p>UNCLASSIFIED</p> <ol style="list-style-type: none"> <li>1. Biophysics</li> <li>2. Circulatory responses</li> <li>3. Acceleration</li> <li>4. Mechanics</li> <li>5. Stress Tests</li> <li>6. Anthropology</li> <li>7. Selection</li> </ol> <p>I. AFSC Project 7222, Task 722202</p> <p>II. Biomedical Labora- tory</p> <p>UNCLASSIFIED</p>
<p>Aerospace Medical Division, 6570th Aerospace Medical Research Laboratories, Wright-Patterson AFB, Ohio Rpt No. AMRL-TDR-62-151. SOME CARDIO- RESPIRATORY RESPONSES OF FLYING AND NON-FLYING PERSONNEL TO DIFFERENT VECTORS OF ACCELERATION WITH CORRE- LATION OF THESE RESPONSES TO OTHER VARIABLES. Final report, Dec 62, iii + 9 pp., incl. illus., tables, 9 refs. Unclassified report</p> <p>The cardiovascular and respiratory responses of test pilots are compared with the AMRL cen- trifuge panel members during headward (+ G<sub>Z</sub>) and forward (+ G<sub>X</sub>) accelerations. Vital capacity decreased in all subjects with increasing for- ward acceleration. No significant</p> <p>( over )</p>	<p>UNCLASSIFIED</p> <ol style="list-style-type: none"> <li>III. Hyde, A. S., Cherniack, N. S., Lindberg, E. F., Capt, USAF, and Whately, D.</li> <li>IV. In ASTIA collection V. Aval fr OTS: \$0.50</li> </ol> <p>UNCLASSIFIED</p>	<p>UNCLASSIFIED</p> <ol style="list-style-type: none"> <li>III. Hyde, A. S., Cherniack, N. S., Lindberg, E. F., Capt, USAF, and Whately, D.</li> <li>IV. In ASTIA collection V. Aval fr OTS: \$0.50</li> </ol> <p>UNCLASSIFIED</p>	<p>UNCLASSIFIED</p> <ol style="list-style-type: none"> <li>III. Hyde, A. S., Cherniack, N. S., Lindberg, E. F., Capt, USAF, and Whately, D.</li> <li>IV. In ASTIA collection V. Aval fr OTS: \$0.50</li> </ol> <p>UNCLASSIFIED</p>